



Superior Livestock Vaccination Programs Designed for Your Management Practices

Management requirements:

- **Please consult your veterinarian or health advisor when selecting the vaccine that qualifies for these programs.**
 - **Always read and follow vaccine label directions.**
 - Always use proper injection sites and techniques. Whenever possible, use Sub Q labeled products.
 - Provide adequate nutrition and minerals to enhance immune system.
 - Include other vaccines or management practices that are unique to your operation or are of value to the calf purchaser.
-

VAC 24

Calves vaccinated on cows at 2 to 4 months of age.

Vaccination requirements:

- 1 dose 7-way, 8-way or 9-way Clostridial.
 - 1 dose viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal & External Parasite Control Recommended.
-

VAC 34

Calves vaccinated on cows at 2-4 weeks prior to shipping.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial. One dose at branding and one dose 2-4 weeks prior to shipping.
 - 1 dose viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal & External Parasite Control Recommended.
-

VAC 34+

Calves vaccinated on cows at branding and 2-4 weeks prior to shipping.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial. One dose at branding and one dose 2-4 weeks prior to shipping.
 - 2 doses viral 5-way (IBR, PI3, BRSV and BVD Type I & II) at branding and 2-4 weeks prior to shipping. IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida 2-4 weeks prior to shipping
 - Internal and external parasite control recommended.
-

VAC 45

Calves vaccinated twice: At branding or 2-4 weeks prior to weaning and booster at weaning, or at weaning and booster to label instructions. Home Raised and Weaned minimum of 45 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial.
 - 2 doses viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida prior to weaning or at weaning and boosted to label instructions.
 - Internal & External Parasite Control Recommended.
-

VAC 45+

Calves vaccinated three times: At branding, 2-4 weeks prior to weaning and booster at weaning or at branding, at weaning and booster to label instructions. Home Raised and Weaned minimum of 45 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial @ branding, prior to or at weaning and boosted to label instructions.
 - 3 doses Modified Live Viral 5 way (IBR, PI3, BRSV, BVD Type I & II) @ branding, prior to weaning or at weaning and boosted to label instructions.
 - 2 doses Mannheimia Haemolytica and/or Pasteurella Multocida @ branding, prior to, or at weaning and boosted to label instructions.
 - Internal & External Parasite Control Required.
-

VAC 60

Calves vaccinated twice: At branding or 2-4 weeks prior to weaning and booster at weaning, or at weaning and booster to label instructions. Home Raised and Weaned minimum of 60 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial.
 - 2 doses viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida prior to weaning or at weaning and boosted to label instructions.
 - Internal & External Parasite Control Recommended.
-

Vac Precon

Vaccinated twice: At arrival and boosted to label instructions, minimum of 14 days prior to delivery. Purchased cattle weaned minimum of 60 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial on arrival and boosted.
 - 2 doses Modified Live Viral 5 way (IBR, PI3, BRSV, BVD Type I & II) on arrival and boosted.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal Parasite Control Recommended, External Parasite Control Recommended.
-